

Pharmacists' private issue dictionary (flags)

Pharmacists have always been able to add to the issues identified by MRM's expert system. In the latest release of MRM (version 3.20) pharmacists can elect to have *new issues* saved to their private dictionary so they can be easily re-called and re-used in other reports. Embedding a pharmacist's frequently used phrases (more commonly known as flags) within the existing workflow of MRM will improve report writing consistency and efficiency.

How it works.

Your private issues dictionary has been seamlessly integrated into your normal workflow.

Adding issues from your dictionary to your report

4 Possible issues identified
0 selected 0 added

Cardiovascular risk reduction
 Mrs Fitzgerald has a history of diabetes and is not taking therapy to reduce risk of cardiovascular events. Treatment with antihyperlipidaemic and antiplatelet agents is recommended in the vast majority of diabetic patients. Confirmation of existing lipid levels, assessment of cardiovascular risk and commencement of appropriate treatment is indicated.

High risk of osteoporosis
 Mrs Fitzgerald is an elderly female with a previous fracture. Investigation of the nature of the fracture (traumatic or not) and

Add New Issue To List +

Next

4 Possible issues identified
0 selected 0 added

Add New Issue

Issue

Issue	Details	New	Close
DIABETES	Although diet control can be used for type 2 diabetes, it is more usual to utilise oral hypoglycaemics and/or metformin. A review of current blood sugar levels is recommended to assess if diet control alone is still appropriate.AAAs	New	Close

Add new issue to personal repository

Cancel
Save

Add new issue

Your private dictionary is automatically searched as you type. Matching flags are listed. Click on an item to copy or [tab] to add a new issue.

Adding / changing your dictionary items

As with MRM identified issues, issues from your dictionary can be customized to the specific needs of the current referral or *Applied every time* which updates the text stored in your dictionary.

4 Possible issues identified
1 selected 1 added

Add New Issue

Issue

Although diet control can be used for type 2 diabetes, it is more usual to utilise oral hypoglycaemics and/or metformin. A review of current blood sugar levels is recommended to assess if diet control alone is still appropriate.

Make changes to my flag

Apply every time issue is detected ?

Cancel Save

Check if you want MRM to update the text in your dictionary.

6 Possible issues identified
0 selected 2 added

Add New Issue

Issue

My new issue text.
I wish to keep this flag for use in other reports

Add new issue to personal repository

Cancel Save

New issues identified in the course of a review can be added to your dictionary for use in other reports

Maintaining your dictionary

A new *Issues Dictionary* management screen added to the Tools menu allows pharmacists to add, edit and delete their items.

MRM Home Referrals Calendar Address Book Archive Statistics Tools My Account MedsCheck

My Dictionary

Search:

Dictionary Entries (flags)

+ Diabetes
Although diet control can be used for type 2 diabetes, it is more usual to utilise oral hypoglycaemics and/or metformin. A review of current blood sugar levels is recommended to assess if diet control alone is still appropriate.AAAs w

+ My new Issue
My new issue text. I wish to keep this flag for use in other reports

+ : Edit phrase (flag)
X : Delete phrase (flag)

Add, edit delete a phrase from your dictionary